SEIZURE ADMINISTRATIVE ORDER

Initiate immediate supportive care:

Protect patient from injury
O2 to maintain sat ≥ 94%
Complete primary and secondary survey as indicated
Vital Signs (FSBG and temperature as indicated)
Cardiac Monitor

Use this Administrative Order on the following patients:

- Seizure activity
- Postictal mental status

Do not use this Administrative Order on the following patients:

Pregnant > 20 weeks gestation and no history of seizures (Follow OB AO)

Actively Seizing

NO

If patient is NOT actively seizing, continue with supportive care

Yes

If patient actively seizing administer

Midazolam (Versed) IM first

- Midazolam (Versed) 0.2mg/kg (max 10 mg) IM
- If IV already established administer *Midazolam* (*Versed*) *0.1mg/kg* (max 5 mg)
- Initiate IV NS/LR TKO
 - If normotensive, consider saline lock with NS flush

Push all meds slowly

- If seizure continues for 10 minutes:
 - No IV access: repeat Midazolam (Versed)
 0.2mg/kg (max 10 mg) IM
 - If IV access: administer *Midazolam (Versed)* 0.1mg/kg (max 5 mg)
- Monitor Airway and breathing

Special Considerations:

If patient is <6 years old, consider febrile seizure and provide cooling techniques

Notification to include:

Seizure Administrative Order, unit number, patient age, gender, and ETA to receiving facility.

Advise if patient is unstable.